

DATE OF APPOINTMENT: _____

OFFICE: _____

Ophthalmic Imaging Request Form

REQUESTING DOCTOR: _____

ADDRESS: _____

DOCTOR PHONE: _____ DOCTOR FAX: _____

PATIENT: _____

DIAGNOSIS: (Necessary) _____



ICD9 CODE: (Necessary) _____

PRIMARY INSURANCE: _____

AUTHORIZATION #: _____

SERVICE REQUESTED (CIRCLE):								
DILATE OU 2.5% MYDFRIN AND 1% MYDRIACYL								
FLUORESCEIN ANGIOGRAM			OPTICAL COHERENCE TOMOGRAPHY			FUNDUS PHOTOS		
PRIMARY EYE:	OD	OS	OD OS			STEREO:	OD	OS
TRANSIT:	OD	OS	INDOCYANINE GREEN ANGIOGRAM (Walnut Creek Office Only)			SEVEN FIELDS:	OD	OS
SWEEP:	OD	OS	OD OS					

PLEASE DRAW AREA OF INTEREST



BARA Outside Imaging Request 9.04

FOR INTERNAL USE ONLY:

VA cc/sc Dil OD OS OU
at _____
OS

Photographer: _____

Report faxed on: _____

Pictures mailed on: _____

**INFORMED CONSENT FOR FLUORESCEIN ANGIOGRAPHY
OR INDOCYANINE GREEN ANGIOGRAPHY**

Patient: _____

- I hereby authorize Bay Area Retina Associates to administer intravenous Fluorescein and/or Indocyanine Green dye for the purpose of performing angiography.
- Angiography is a diagnostic procedure in which a rapid sequence of photographs are taken to document the blood circulation of the retina/choroid. The dye is usually injected into a vein in the arm, forearm or hand.
- Since the fluorescein dye is a very bright yellow, the skin may appear jaundiced for a few hours and then the yellow color disappears. The dye is excreted through the kidney causing the urine to be a bright yellow for 24-36 hours.
- Indocyanine Green is excreted through the bile and will show in the stool as greenish. The coloration of these dyes are considered to be a normal result of the after effects.

Patients with known allergies to shellfish or iodine should not have ICG Angiography.

- Less than 0.2% of all patients to whom these tests are administered have any of the listed side effects. Documented adverse reactions to the dyes which can occur include: nausea, headache, upset stomach, vomiting, light-headedness, fainting, hives or itching. Even more rarely, severe allergic reactions (anaphylaxis) can occur and be life threatening. Extravasation of the fluorescein dye out of the blood vessel is painful and every effort is made to prevent this from occurring.

FOR WOMEN:

- Intravenous fluorescein/ICG is usually not administered to pregnant women, although there is no scientific evidence to suggest that it might harm unborn babies.
- _____ To the best of my knowledge, I am not pregnant.
- I hereby authorize and direct Bay Area Retina Associates and/or their designees to provide such additional services as they may deem necessary and reasonable.
- I understand that no guarantees of any kind regarding these procedures have been made to me.
- I consent to the use of the above photographs and other materials for scientific purposes, provided my identity is not revealed by the pictures or the descriptive text accompanying them.

Patient (or person authorized to sign for patient)

Date

Witness

Date